

THE MARYLAND ENTOMOLOGIST

Insect and related-arthropod studies in the Mid-Atlantic region



Volume 7, Number 3

September 2019

MARYLAND ENTOMOLOGICAL SOCIETY
www.mdentsoc.org

Executive Committee:	President	Frederick Paras
	Vice President	Philip J. Kean
	Secretary	Janet A. Lydon
	Treasurer	Edgar A. Cohen, Jr.
	Historian	(vacant)
	Journal Editor	Eugene J. Scarpulla
	E-newsletter Editor	Aditi Dubey

The Maryland Entomological Society (MES) was founded in November 1971, to promote the science of entomology in all its sub-disciplines; to provide a common meeting venue for professional and amateur entomologists residing in Maryland, the District of Columbia, and nearby areas; to issue a periodical and other publications dealing with entomology; and to facilitate the exchange of ideas and information through its meetings and publications. The MES was incorporated in April 1982 and is a 501(c)(3) non-profit, scientific organization.

The MES logo features an illustration of *Euphydryas phaëton* (Drury) (Lepidoptera: Nymphalidae), the Baltimore Checkerspot, with its generic name above and its specific epithet below (both in capital letters), all on a pale green field; all these are within a yellow ring double-bordered by red, bearing the message “● Maryland Entomological Society ● 1971 ●”. All of this is positioned above the Shield of the State of Maryland. In 1973, the Baltimore Checkerspot was named the official insect of the State of Maryland through the efforts of many MES members.

Membership in the MES is open to all persons interested in the study of entomology. All members receive the annual journal, *The Maryland Entomologist*, and the monthly e-newsletter, *Phaëton*. Institutions may subscribe to *The Maryland Entomologist* but may not become members.

Annual Dues:	Individual Membership	\$10.00
	Household Membership	\$15.00
	Junior Membership (full-time student)	\$5.00
	Institutional Subscription	\$10.00

Prospective members should send to the Treasurer full dues for the current MES year (October – September), along with their full name, address, telephone number, e-mail address, and entomological interests. Applications can be downloaded from the MES website: www.mdentsoc.org. Send remittances, payable to the Maryland Entomological Society, and any address changes to the Treasurer: Edgar J. Cohen, Jr., 5454 Marsh Hawk Way, Columbia, MD 21045-2246, edcohenfam@yahoo.com.

Back issues of *The Maryland Entomologist* and recent issues of the *Phaëton* are available to members, via the Journal Editor, Eugene J. Scarpulla, ejscarp@comcast.net. Please contact the Journal Editor for availability and cost.

Meetings are held on the third Friday of October, November, February, March, April and May at 7:00 p.m. in Room 4 of the Biological Sciences Building, University of Maryland Baltimore County (UMBC), or occasionally at another announced site.

Past Presidents of the MES

1971-1973	Austin P. Platt	1984-1985	Charles L. Staines, Jr.
1973-1974	Ronald W. Hodges	1985-1986	Thomas E. Wallenmaier
1974-1975	Douglas C. Ferguson	1986-1987	Eugene J. Gerberg
1975	Raymond B. Nagle	1987-1988	Austin P. Platt
1975-1977	William A. Andersen	1988-1989	Philip J. Kean
1977-1978	Robert T. Mitchell	1989-1990	Nathan Erwin
1978-1979	Elaine R. Hodges	1990-1991	Stephen J. Harrison
1979-1980	Richard H. Smith, Jr.	1991-1992	Thomas E. Wallenmaier
1980-1981	Timothy P. Karpetsky	1992-1993	Mary C. Fenton
1981-1982	John F. Carroll	1993-2013	Frederick Paras
1982-1983	Theodore L. Bissell	2013-2015	Timothy Foard & Frederick Paras
1983-1984	Robin G. Todd	2015-present	Frederick Paras

Editor's Note

The Maryland Entomologist is regional in scope. This year's issue features studies conducted in Maryland, Virginia, West Virginia, and the District of Columbia. This year's authors hail from six jurisdictions: one from Maine, one from Maryland, one from Oregon, one from Virginia, two from West Virginia, and two from the District of Columbia.

Frank G. Guarnieri reviewed incidental captures of *Calosoma externum* (Say) (Coleoptera: Carabidae) on Maryland's Coastal Plain during a lepidopteran monitoring and control program. **Brent W. Steury** documented the beetles (Coleoptera) associated with flowers of lizard's Tail, *Saururus cernuus* L. (Saururaceae), in Calvert County, Maryland. **Gary D. Ouellette** investigated the diversity of Sciapodinae (Diptera: Dolichopodidae) in the Greater Washington, District of Columbia area. **Mark J. Hepner** and **Ernest W. Smith** established recent records of the Yellow-banded Bumble Bee, *Bombus terricola* Kirby (Hymenoptera: Apidae), in West Virginia. **Jonathan R. Mawdsley**, **Davia M. Palmeri**, and **Mark Humpert** documented the occurrence of bumble bees, *Bombus* Latreille (Hymenoptera: Apidae), in State Wildlife Action Plans as new opportunities for conservation.

Call for Potential Authors

Dig into your field notebooks and consider publishing in *The Maryland Entomologist*. [*Don't wait till you croak; it's too late then!*] I am sure there are much data out there just waiting to be published. Put your findings in print for the world to see. Remember, your study benefits no one if you are the only person with access to it. I look forward to an inbox overflowing with submittals for the 2020 issue. Please e-mail first drafts to me by April 1, 2020. Thanks for your consideration.

Eugene J. Scarpulla
Editor

Incidental Capture of *Calosoma externum* (Say) (Coleoptera: Carabidae) at Various Sites across the Coastal Plain of Maryland between 1988 and 1993 during a Monitoring and Control Program Targeting Lepidopteran Agricultural Pests

Frank G. Guarnieri

39 Roosevelt Avenue, Waterville, Maine 04901; fgguarnieri@aol.com

Abstract: *Calosoma externum* (Say) (Coleoptera: Carabidae) is a large (24–35 mm) and attractive beetle that is not regularly encountered in the Mid-Atlantic region. Yet, *C. externum* was found in five counties on the Coastal Plain of Maryland between 1988 and 1993. The beetles were bycatch in ultraviolet light traps during an integrated pest management program targeting various species of destructive moths. Published collecting records of *C. externum* in and around Maryland are reviewed. This species may be more widespread in the Mid-Atlantic region than is currently recognized. This paper does not present enough data to make any conclusions regarding population numbers. Photographs are presented comparing *C. externum* and two similar carabid species with which it might be confused in Maryland.

Calosoma externum (Say) (Coleoptera: Carabidae) is a large (24–35 mm [0.9–1.4 in]) and attractive beetle (Figure 1). The dorsal surface is polished black. The pronotum and elytra have brilliant metallic blue or purple margins. Typically, *Calosoma* species have broad elytra that, in dorsal view, give the beetles a somewhat blocky appearance. In comparison, the elytra of *C. externum* are long and narrow. The result is an elongate and streamlined habitus, hence its common name, the Narrow Searcher. An excellent photomontage depicting all Mid-Atlantic species of *Calosoma* can be seen in Evans (2009).

Calosoma species are called searcher or caterpillar hunter beetles. Larvae and adults are well known to prey upon many insect varieties, lepidopteran larvae in particular. *Calosoma externum* has been noted to consume many agricultural pest species of Orthoptera, Coleoptera, and Lepidoptera, notably caterpillars of: *Mythimna unipuncta* (Haworth) (Lepidoptera: Noctuidae), the Armyworm; *Malacosoma americanum* (Fabricius) (Lepidoptera: Lasiocampidae), the Eastern Tent Caterpillar; and *Lymantria dispar* (Linnaeus) (Lepidoptera: Erebiidae), the Gypsy Moth (Burgess and Collins 1917).

Based upon Burgess and Collins (1917), Gidaspow (1959), and Bousquet (2012), the range of *Calosoma externum* extends across eastern North America from Massachusetts to Georgia in the East, and Minnesota and Nebraska to Texas in the West. Burgess and Collins (1917) and Gidaspow (1959) report this species is more common in the “southern” part of its range. Bousquet (2012) reports that isolated records from Ontario and Vermont are probably “strays.” According to Larochelle and Larivière (2003), *C.*

externum is most likely to be found in “cultivated fields, pastures, vacant lots, and open forests.”

Despite three decades of field experience in the Mid-Atlantic region, I have never encountered *Calosoma externum* in the wild. Based on anecdotal reports and casual discussions with professional and avocational carabid specialists, this beetle is most likely to be found in the South-Central United States. (The type specimen described by Thomas Say in 1823 was collected in Arkansas [Burgess and Collins 1917].) Such speculation is supported by a small data set (ten records between 1972 and 2017) on BugGuide (2019). The northernmost records of *C. externum* on the website are from Indianapolis, Indiana and Lincoln, Nebraska. There are two records from Missouri and single records each from Illinois, Kansas, Kentucky, North Carolina, Oklahoma, and Texas.



Figure 1. *Calosoma externum* (Say). **A) Male:** Maryland, Dorchester County, Church Creek, 15 July 1988, collected at ultraviolet light, 30 mm (1.2 in) (MDAG collection). Males have enlarged protarsal segments compared to females. The bright blue or purple margins are difficult to reproduce in photographs but can be seen fairly well in this specimen. **B) Female:** Maryland, Worcester County, Indiantown, 19 July 1993, collected at ultraviolet light, 33 mm (1.3 in) (MDAG collection).

Regarding the presence of *Calosoma externum* in and around Maryland, Burgess and Collins (1917) found eighteen specimens under electric lights in the District of Columbia between 1909 and 1912. Subsequent published records become elusive. In recent decades, many detailed surveys of carabid beetles were conducted in the District of Columbia, Maryland, southern Pennsylvania, and northern Virginia (Erwin 1981; Stork 1984; Bailey et al. 1994; Anderson et al. 1995; Evans 2008; Kim and Piechnik 2009; Guarnieri 2010; Staines and Staines 2011; Fritzler and Strazanac 2012; Steury and Messer 2014, 2015, 2017; Steury et al. 2014). Most recorded at least one *Calosoma* species, but only one survey (Steury and Messer 2017) reported *C. externum* (Calvert County, Cove Point beach, 26 September, one specimen in beach drift). Steury (in litt.) reported that the specimen was dead and missing an entire elytron. It is uncertain precisely where the specimen may have originated before floating onshore.

Recently, I encountered six Maryland specimens of *Calosoma externum* that were collected between 1988 and 1993. Four were located at the Maryland Department of Agriculture in Annapolis, Maryland (MDAG): Charles Co., Hughesville, 25 May 1993; Dorchester Co., Church Creek, 15 July 1988; Somerset Co., Princess Anne, 21 June 1988; and Worcester Co., Indiantown, 19 July 1993. Two were at the Carnegie Museum of Natural History in Pittsburgh, Pennsylvania (CMNH): Cecil Co., 12 August 1993 and Dorchester Co., Rhodesdale, 17 July 1993.

According to Gaye L. Williams (Entomologist, Plant Protection and Weed Management, MDAG) (in litt.), these six beetles were all likely long forgotten bycatch of an integrated pest management program that was started by the University of Maryland and taken over by the MDAG. The study involved numerous collaborators including the CMNH. Farmers and field scouts operated ultraviolet light traps and sent in daily catches. The samples were examined for 16 lepidopteran crop pests including: *Agrotis ipsilon* (Hufnagel) (Lepidoptera: Noctuidae), the Black Cutworm; *Helicoverpa zea* (Boddie) (Lepidoptera: Noctuidae), the Corn Earworm; and *Ostrinia nubilalis* (Hübner) (Lepidoptera: Crambidae), the European Corn Borer. Farmers used the data to coordinate control applications based on population surges of pest species.

In addition to the two Maryland specimens discussed above, the CMNH contained over 100 specimens of *Calosoma externum*. The data support the notion of a historically widespread eastern species that now appears to be most common in the South-Central United States. There are scattered records from the late 1800s and early 1900s from the District of Columbia, Maryland, Michigan, New York (restricted to Long Island), and Pennsylvania. There are two specimens from Virginia, New Kent Co., 31 December 1952. Yet, the vast majority of records, including many recently collected specimens, are from Oklahoma, Tennessee, and Texas.

The data from the MDAG survey suggest that *Calosoma externum* may be more widespread over the Coastal Plain of Maryland than is currently appreciated. But it remains unclear how these large and distinctive beetles generally go unnoticed. They may be uncommon or simply secretive. According to Robert L. Davidson (Collection Manager, Division of Invertebrate Zoology, CMNH) (in litt.), even if this species

regularly occurs in large corn and soybean fields, it could escape detection as these habitats are not typically sampled in carabid surveys.

Identification of *Calosoma externum* is fairly straightforward. However, in my experience, there are two common related species on the Coastal Plain of Maryland that might cause confusion: *Carabus sylvosus* Say (Carabidae), Woodland Ground Beetle (Figure 2A) and *Calosoma sayi* Dejean (Carabidae), Black Caterpillar Hunter, (Figure 2B).



Figure 2. *Carabus sylvosus* Say and *Calosoma sayi* Dejean. A) *Carabus sylvosus* female: the basal angles of the pronotum are highly reflexed (black arrow) and the elytral striae are indistinct. Maryland, Worcester County, Pocomoke River State Park (38.1355°, -75.4384°), 20 July 2000, crawling on the ground in mature woods at night, 30 mm (1.2 in) (Frank G. Guarnieri collection). B) *Calosoma sayi* male: the elytral striae are interrupted by small metallic red, yellow, or green punctures. Maryland, Worcester County, Snow Hill (38.1879°, -75.3769°), 17 July 2000, in a parking lot at night under bright lights, 29 mm (1.1 in) (Frank G. Guarnieri collection).

Carabus sylvosus: Burgess and Collins (1917) and Ciegler (2000) also note that *Calosoma externum* and *Carabus sylvosus* are quite similar in appearance. In the latter species, the basal angles of the pronotum are more strongly reflexed and the elytral striae are indistinct. *Carabus sylvosus* is fairly easy to find running on the ground in open woods at night. The *Carabus sylvosus* specimen depicted in Figure 2A was found ~8 km (~5 mi) from the approximate location where the *Calosoma externum* specimen shown in Figure 1B was collected.

Calosoma sayi: Males, in particular, of *C. sayi* tend to have elytra that are nearly parallel-sided. The result is a thinner habitus that is closer to *C. externum* than the other members of *Calosoma* that have much wider abdomens (see the figure in Evans [2009], however, note that the *C. sayi* in the photomontage is a female specimen). *Calosoma sayi* can be identified by rows of small metallic red, yellow, or green punctures on the elytra. This species is readily attracted to light and can be abundant at night in 24-hour gas station parking lots in open agricultural areas. The *C. sayi* specimen shown in Figure 2B was found ~3 km (~2 mi) from the approximate location where the *C. externum* specimen shown in Figure 1B was collected.

To conclude, it is important to note that the Maryland records of *Calosoma externum* discussed in this report represent an extremely small data set. Thus, this paper does not present enough data to make conclusions regarding population numbers or trends. A comprehensive survey of the major regional insect collections to look for additional modern Mid-Atlantic records would be most useful. With a larger data set, it may be possible to show these beetles occur in “boom and bust” cycles, with the majority of individuals being collected in just a few years surrounded by long periods of relative scarcity. Such a pattern may, in part, explain the general paucity of observations in the field. Furthermore, it would be interesting to determine if *C. externum* numbers wax and wane in conjunction with outbreaks of particular agricultural pests or if there are any effects from pesticide applications. But for now, the intent herein is only to make readers aware of this interesting species and be vigilant to its possible occurrence, particularly in lowland agricultural fields, throughout the Mid-Atlantic region.

ACKNOWLEDGMENTS

I am indebted to Robert L. Davidson (CMNH) and Gaye L. Williams (MDAG) for very useful discussions and also for giving me access to the collections under their care. Two anonymous reviewers made most constructive comments including excellent ideas regarding possible directions of future research.

LITERATURE CITED

Anderson, J.M., J.C. Mitchell, A.A. Hall, and R.L. Hoffman. 1995. Ground Beetles (Coleoptera: Carabidae) from Quantico Marine Corps Base, Virginia. *Banisteria* 6:3–16.

- Bailey, R.R., E.P. van den Berghe, and B.C. McCarthy. 1994. New records and activity patterns of carabid beetles (Coleoptera: Carabidae) in Western Maryland. *The Coleopterists Bulletin* 48(4):319–323.
- Bousquet, Y. 2012. Catalogue of Geadephaga (Coleoptera, Adephaga) of America, north of Mexico. *ZooKeys* 245:1–1722.
- BugGuide. 2019. Species *Calosoma externum* – Narrow Searcher. Hosted by Iowa State University Department of Entomology. Available at: <https://bugguide.net/node/view/3880/data>. Last accessed on 28 March 2019.
- Burgess, A.F., and C.W. Collins. 1917. The genus *Calosoma*: including studies of seasonal histories, habits, and economic importance of American species North of Mexico and of several introduced species. *United States Department of Agriculture Bulletin* 417:1–124.
- Ciegler, J.C. 2000. *Ground Beetles and Wrinkled Bark Beetles of South Carolina (Coleoptera: Geadephaga: Carabidae and Rhysodidae)*. Biota of South Carolina, Volume 1. South Carolina Agriculture and Forestry Research System, Clemson University, Clemson, SC. 149 pp.
- Erwin, T.L. 1981. Natural history of Plummers Island, Maryland. XXVI. The ground beetles of a temperate forest site (Coleoptera: Carabidae): an analysis of fauna in relation to size, habitat selection, vagility, seasonality, and extinction. *Bulletin of the Biological Society of Washington*, 5:104–224.
- Evans, A.V. 2008. The 2006 Potomac Gorge BioBlitz: Beetles. *Banisteria* 32:44–51.
- Evans, A.V. 2009. The forest caterpillar hunter, *Calosoma sycophanta*, an Old World species confirmed as part of the Virginia beetle fauna (Coleoptera: Carabidae). *Banisteria* 34:33–37.
- Fritzler, C.J., and J.S. Strazanac. 2012. *Survey of Ground Beetles (Carabidae) and Other Select Coleoptera (Scarabaeidae, Geotrupidae, Trogidae, Tenebrionidae, Silphidae) in Catoctin Mountain Park*. Report to the National Park Service, United States Department of the Interior. 101 pp.
- Gidaspow, T. 1959. North American caterpillar hunters of the genera *Calosoma* and *Callisthenes* (Coleoptera, Carabidae). *Bulletin of the American Museum of Natural History* 116(3):225–344.
- Guarnieri, F.G. 2010. A survey of the beetles (Coleoptera) of Pocomoke River State Park, Worcester County and Tuckahoe State Park, Caroline County, Maryland. *The Maryland Entomologist* 5(2):5–28.
- Kim, K.C., and D.A. Piechnik. 2009. *Ecological Analysis of Arthropod Diversity and Development of a Monitoring Plan for Gettysburg National Military Park and*

- Eisenhower National Historic Site*. Technical Report NPS/NER/NRTR—2009/143. National Park Service, Philadelphia, PA. 213 pp.
- Larochelle, A., and M.-C. Larivière. 2003. *A Natural History of the Ground-Beetles (Coleoptera: Carabidae) of America North of Mexico*. Pensoft Series Faunistica No. 27. Pensoft Publishers, Sofia, Bulgaria-Moscow, Russia. 583 pp.
- Staines, C.L., and S.L. Staines. 2011. The Carabidae (Insecta: Coleoptera) of Eastern Neck National Wildlife Refuge, Maryland. *Banisteria* 38:71–84.
- Steury, B.W., and P.W. Messer. 2014. Twelve ground beetles new to Virginia or the District of Columbia and an annotated checklist of the Geadephaga (Coleoptera, Adephaga) from the George Washington Memorial Parkway. *Banisteria* 43:40–55.
- Steury, B.W., and P.W. Messer. 2015. Noteworthy beetle records from Virginia and Maryland (Coleoptera: Anthicidae, Buprestidae, Carabidae). *Banisteria* 45:61–62.
- Steury, B.W., and P.W. Messer. 2017. Three carabid beetles (Coleoptera: Carabidae) new to Maryland and a preliminary annotated checklist for Cove Point, Calvert County, Maryland. *The Maryland Entomologist* 7(1):35–47.
- Steury, B.W., P.W. Messer, and J.F. Cavey. 2014. Noteworthy beetle records from Virginia, Maryland, and the District of Columbia (Coleoptera: Carabidae and Chrysomelidae). *Banisteria* 44:23–25.
- Stork, N.E. 1984. Additions to the list of Carabidae (Coleoptera) in the fauna of Plummers Island, Maryland. *The Coleopterists Bulletin* 38(2):137–141.

Beetles (Coleoptera) Associated with Flowers of Lizard's Tail, *Saururus cernuus* L. (Saururaceae), in Calvert County, Maryland

Brent W. Steury

8316 Woodacre Street, Alexandria, Virginia, 22308; bsteury@cox.net

Abstract: Beetles associated with flowers of lizard's tail, *Saururus cernuus* L. (Saururaceae), are documented from a freshwater swamp in southern Maryland. Six hundred forty-six beetles representing 29 species in 14 families were documented on flowers of *S. cernuus* during eight hours of collection effort over four days in June. Mating was observed in three species of Cerambycidae, indicating a possible connection between pollen consumption and copulation. Pollen and other flower parts may be food sources for some beetles. The most common beetle on flowers of *S. cernuus* was *Isomira sericea* (Say) (Tenebrionidae). The most species-rich family was Mordellidae. Other arthropod groups observed on flowers of *S. cernuus* included: Arachnida (Araneae and Acari), Collembola, Diptera, Hemiptera, Hymenoptera, Lepidoptera, and Orthoptera.

INTRODUCTION

Lizard's tail, *Saururus cernuus* L. (Saururaceae), is a perennial, rhizomatous, herbaceous plant common on hydric soils, typically in shady habitats. It is common throughout the Coastal Plain and Piedmont of Maryland and Virginia (Brown and Brown 1984; Virginia Botanical Associates 2018) and occurs from Canada to Florida, westward to Texas (Buddell and Thieret 1997). The lizard's tail family (Saururaceae) is a small, primitive, plant family containing only six species in the order Piperales (Raju 1961). Only *S. cernuus* and yerba mansa, *Anemopsis californica* (Nutt.) Hook. & Arn., are native to North America; the others are Asian (Buddell and Thieret 1997). *Saururus cernuus* reaches heights of just over 1 m (3.3 ft) and has entire, cordate leaves, the lower ones the largest, generally 15 cm (9 in) long and 10 cm (4 in) wide. Each plant produces one or two terminal, crook-shaped racemes on each branch. The showy, bright white, aromatic inflorescences may be over 20 cm (8 in) long and 1 cm (0.4 in) wide and contain up to 350 small, tightly aggregated flowers with 6 or 8 protruding stamens in two whorls or 3 in a single whorl (Raju 1961). Anthesis proceeds from the base of the racemes to the tip, rendering a gradient of flowering and fruit maturation and a straightening of the inflorescence. In the southern United States, anthesis proceeds at approximately 1.52 cm (0.6 in) per day (Thien et al. 1994). Thien et al. (1994) concluded pollination was primarily insect-mediated or wind-dependent. *Saururus cernuus* propagates both vegetatively and by seed; however, vegetative reproduction may be more important to population persistence since seedlings have not been reported from field studies (Hall 1940, Penfound et al. 1945). Thien et al. (1994) attempted to discern the pollination mode for *S. cernuus* in Louisiana based on observations of potential pollinators and observed

Diptera and Coleoptera feeding on pollen. This study documents the species of Coleoptera and other arthropod orders associated with flowers of a large population of *S. cernuus* in Maryland.

STUDY SITE

The study site at Cove Point, in Calvert County, Maryland, is located at 38°23'30.6", -76°24'02.2" in a forested swamp with a canopy of red maple, *Acer rubrum* L. (Sapindaceae), and blackgum, *Nyssa sylvatica* Marshall (Cornaceae). The herb layer is strongly dominated by a crescent-shaped patch of *S. cernuus* (Figure 1) with a maximum length and width of approximately 70 m (230 ft) by 20 m (66 ft). The site is situated near the Chesapeake Bay and is bordered by a non-tidal, freshwater marsh known for its high number of state rare plant species (Steury 1999).



Figure 1: The study site at Cove Point, Calvert County, Maryland on 25 June 2018, showing lizard's tail (*Saururus cernuus* L.) in foreground, under a canopy of red maple (*Acer rubrum* L.) and blackgum (*Nyssa sylvatica* Marshall).

MATERIALS and METHODS

The top was cut from a plastic, 3.8 L (1 gal) milk jug and a thin film of water was added. The jug was held under an *S. cernuus* inflorescence as it was bent into the jug and shaken. The process was conducted for the first 30 minutes of each hour, from 9:00 a.m. to 12:30 p.m. on 21–23 June 2018, and from 11:00 a.m. to 2:30 p.m. on 25 June 2018. The process did not damage the inflorescences other than the abscission of some dehiscent anthers from the flowers. Beetles fell to the bottom of the jug and were held by the surface tension of the water. They were removed from the jug by hand and placed in a glass vial containing 95% ethanol. One male and one female (when available) of each species was pinned, labeled, and deposited in the entomology collection at the United States National Museum, Smithsonian Institution, Washington, DC. No observed beetle species eluded capture, and it was noted that inflorescences which on close inspection appeared to contain no beetles, often yielded Coleoptera in the bottom of the jug. Individuals of other arthropod orders captured during the survey were released after being recorded in a field notebook. During each day of the survey, *S. cernuus* flowers at various stages of anthesis were present. No leaves of *S. cernuus* were inserted in the jug and arthropods observed on leaves were not included in the tally. Weather was recorded on 21 June as partly cloudy with a high temperature of 28.9 °C (84 °F) during the survey; 22 June as intermittent light rain and a high temperature of 23.9 °C (75 °F); 23 June as overcast and humid with a high temperature of 27.8 °C (82 °F); and 25 June as mostly sunny with a high temperature of 28.3 °C (83 °F).

RESULTS and DISCUSSION

A total of 646 beetles of 29 species in 14 families were captured from flowers of *S. cernuus* at Cove Point Marsh during eight hours of sampling over four days (Table 1). The most common beetle captured was *Isomira sericea* (Say) (Tenebrionidae) (n = 347). Other common species were *Mordellistena masoni* Liljeblad (n = 146), *Glipostenoda ambusta* (LeConte) (n = 21), *Mordella marginata* Melsheimer (n = 20), and *Falsomordellistena pubescens* (Fabricius) and *Mordellistena liturata* (Melsheimer) (n = 18 each) (all Mordellidae). The families with the highest species richness were Mordellidae (n = 10 species), Cerambycidae (n = 4 species), and Curculionidae (n = 3 species). Families with the highest number of individuals were Tenebrionidae (n = 347), Mordellidae (n = 233), and Cerambycidae (n = 27). Three of four species of Cerambycidae: *Strangalia luteicornis* (Fabricius), *Typocerus lugubris* (Say), and *T. velutinus velutinus* (Olivier), were observed in copula on inflorescences of *S. cernuus*, with the female apparently feeding on pollen during copulation, indicating a possible connection between pollen consumption and mating. One uncaptured and unidentified mordellid that was observed through a 10× hand lens appeared to be feeding on the base of a *S. cernuus* flower ovary, creating a dark scar when finished. Larger beetles (e.g., Cerambycidae and Cantharidae) were observed clinging to the stamens on the exterior edge of the racemes, while smaller beetles (e.g., Mordellidae and *Isomira sericea*) were primarily observed inside the inflorescence along the rachis. The largest beetle captured on flowers of *S. cernuus* was *T. v. velutinus* (17 mm [0.7 in]) and the smallest was *Orthoperus glaber* (LeConte) (Corylophidae) (0.6 mm [0.02 in]). The record of *Lebia*

ornata Say (Carabidae) is the first for Cove Point, increasing the known carabid fauna from the site to 70 species (Steury and Messer 2017).

Based on findings of this study, *S. cernuus* is an important food source for many species of beetles in at least 14 families that may feed upon its pollen and other flower parts. Further study may reveal additional beetle families associated with *S. cernuus*. Other arthropod groups observed on flowers of *S. cernuus* included: Arachnida (Araneae: one spider with a captured wasp and one spider with a captured syrphid fly; and Acari: mites on some beetles), Collembola, Diptera, Hemiptera, Hymenoptera (wasps and ants only), Lepidoptera (larvae, and one adult, dark morph female, Eastern Tiger Swallowtail [*Papilio glaucus* Linnaeus (Papilionidae)]), and Orthoptera. The importance of the large, showy, aromatic inflorescences to a plant that almost exclusively reproduces vegetatively remains unknown.

Table 1: Number of beetles captured on flowers of *Saururus cernuus* L. at Cove Point, Calvert County, Maryland, during two hours of daily search effort on 21–23 June and 25 June 2018. Taxa are listed alphabetically by family, genus, and species.

Family, Genus, Species, and Author	21 June	22 June	23 June	25 June	Total Captured
Anthicidae (antlike flower beetles)					
<i>Macratria murina</i> (Fabricius)	7			3	10
Cantharidae (soldier beetles)					
<i>Chauliognathus marginatus</i> (Fabricius)	4				4
<i>Rhagonycha angulata</i> (Say)	1				1
Carabidae (ground beetles)					
<i>Lebia ornata</i> Say		1			1
Cerambycidae (long-horned beetles)					
<i>Obrium rufulum</i> Gahan	1				1
<i>Strangalia luteicornis</i> (Fabricius)	2	1	3	2	8
<i>Typocerus lugubris</i> (Say)	3				3
<i>Typocerus velutinus velutinus</i> (Olivier)	3		6	6	15
Chrysomelidae (leaf beetles)					
<i>Diabrotica undecimpunctata howardi</i> Barber	3	4	2	2	11
Corylophidae (minute hooded beetles)					
<i>Orthoperus glaber</i> (LeConte)	1			1	2
Curculionidae (true weevils)					
<i>Anthonomus signatus</i> Say			1		1
<i>Geraeus picumnus</i> (Herbst)			1		1
<i>Tyloderma variegatum</i> (Horn)				1	1

Family, Genus, Species, and Author	21 June	22 June	23 June	25 June	Total Captured
Elateridae (click beetles)					
<i>Glyphonyx nanus</i> Smith and Balsbaugh				1	1
Latridiidae (Minute Brown Scavenger Beetles)					
<i>Melanophthalma</i> Motschulsky sp.				1	1
Melyridae (soft-winged flower beetles)					
<i>Hypebaeus oblitus</i> (LeConte)	1				1
Mordellidae (tumbling flower beetles)					
<i>Falsomordellistena hebraica</i> (LeConte)	1				1
<i>Falsomordellistena pubescens</i> (Fabricius)	9	1	2	6	18
<i>Glipostenoda ambusta</i> (LeConte)	6	5	4	6	21
<i>Mordella marginata</i> Melsheimer	5	9	3	3	20
<i>Mordella obliqua</i> LeConte	1				1
<i>Mordellistena fuscata</i> (Melsheimer)			1	1	2
<i>Mordellistena liturata</i> (Melsheimer)	3	3	5	7	18
<i>Mordellistena masoni</i> Liljeblad	54	19	46	27	146
<i>Mordellistena vapida</i> LeConte	1			1	2
<i>Mordellistena vera</i> Liljeblad	2	1		1	4
Scarabaeidae (scarab beetles)					
<i>Valgus canaliculatus</i> (Olivier)	1				1
Scraptiidae (false flower beetles)					
<i>Allopoda lutea</i> (Haldeman)	2			1	3
Tenebrionidae (darkling beetles)					
<i>Isomira sericea</i> (Say)	106	64	82	95	347
Total	217	108	156	165	646

ACKNOWLEDGMENTS

Appreciation is extended to the Cove Point Natural Heritage Trust for providing funding for the study. Ted C. MacRae (Senior Research Entomologist, Monsanto Company, St. Louis, Missouri) and Eugene J. Scarpulla provided helpful reviews of the manuscript.

LITERATURE CITED

Brown, M.L., and R.G. Brown. 1984. *Herbaceous Plants of Maryland*. Port City Press, Inc., Baltimore, MD. 1127 pp.

Buddell, G.F., II, and J.W. Thieret. 1997. Saururaceae. Pages 37–38 in: *Flora of North America* Editorial Committee (Editors). *Flora of North America North of Mexico Volume Three: Magnoliophyta: Magnoliidae and Hamamelidae*. Oxford University Press, Oxford, UK. 616 pp.

- Hall, T.F. 1940. The biology of *Saururus cernuus* L. *The American Midland Naturalist* 24(1):253–260.
- Penfound, W.T., T.F. Hall, and A.D. Hess. 1945. The spring phenology of plants in and around the reservoirs of north Alabama with particular reference to malaria control. *Ecology* 26(4):332–352.
- Raju, M.V.S. 1961. Morphology and anatomy of the Saururaceae. I. Floral anatomy and embryology. *Annals of the Missouri Botanical Garden* 48(2):107–124.
- Steury, B.W. 1999. Annotated list of vascular plants from a nontidal barrier wetland along the Chesapeake Bay in Calvert County, Maryland. *Castanea* 64(2):187–200.
- Steury, B.W., and P.W. Messer. 2017. Three carabid beetles (Coleoptera: Carabidae) new to Maryland and a preliminary annotated checklist for Cove Point, Calvert County, Maryland. *The Maryland Entomologist* 7(1):35–47.
- Thien, L.B., E.G. Ellgaard, M.S. Devall, S.E. Ellgaard, and P.F. Ramp. 1994. Population structure and reproductive biology of *Saururus cernuus* L. (Saururaceae). *Plant Species Biology* 9(1):47–55.
- Virginia Botanical Associates. 2018. Digital Atlas of the Virginia Flora. Virginia Botanical Associates, Blacksburg, VA. Available at: <http://www.vaplantatlas.org>. Accessed: 23 June 2018.

Diversity of Sciapodinae (Diptera: Dolichopodidae) in the Greater Washington, District of Columbia Area

Gary D. Ouellette

5015 Nantucket Road, College Park, Maryland 20740; gary.d.ouellette@gmail.com

Abstract: Faunal inventories are critical for the conservation of biological resources and diversity. At present, detailed knowledge of Sciapodinae (Diptera: Dolichopodidae) distributions in the Nearctic region is still poorly understood. Herein, I investigate the composition of sciapodine flies in the greater Washington, District of Columbia area. A total of 1657 specimens were identified from the collections of the National Museum of Natural History and the author, representing four genera and 19 species. The most common species, by total occurrences, were *Condylostylus caudatus* (Wiedemann), *C. patibulatus* (Say), and *C. sipho* (Say), which accounted for nearly 83% of all identified flies. The species *Amblypsilopus dorsalis* (Loew), *Condylostylus banksii* (Van Duzee), and *Mesorhaga albiciliata* (Aldrich) are reported for the first time in Maryland. Results presented in this study serve as a baseline for future ecological and biodiversity comparison.

INTRODUCTION

Detailed knowledge of the biota of any site or region is critical for community characterization and environmental impact studies, and is an initial step for biological and natural resource conservation planning (Longino and Colwell 1997, Hughes et al. 2000, Green et al. 2009). Since the early 20th century, there has been considerable interest in the biodiversity of the greater Washington, District of Columbia (DC) area, much of it investigated by members of the Washington Biologists' Field Club (Banks et al. 1916, Alexander and McAtee 1920, McAtee 1921, Malloch et al. 1931, Perry 2007). Although there are more recent studies focused on the biological diversity of insects from the Potomac Gorge (Mathis and Foster 2007, Brown and Bahr 2008, Evans 2008), inventories of the biota of the greater Washington, DC area are still limited for many taxonomic groups.

The Sciapodinae are one of the 14 recognized Nearctic subfamilies in the family Dolichopodidae (Insecta: Diptera) and are comprised of metallic-colored flies usually characterized by a deeply excavated vertex and a widely diverging branch of vein M (Robinson 1964). In the Nearctic region north of Mexico, the subfamily is represented by four genera and 71 species (Pollet et al. 2004) with adults abundant in sunny and moist habitats (Bickel 1994). Generally, adults and larvae of Dolichopodidae are predaceous, occupying an important position in most terrestrial ecosystems. In addition, members of the family have shown promise as biological indicators of environmental quality for conservation purposes (Pollet and Grootaert 1991, 1996; Pollet 2001; Pollet et al. 2003).

Despite these important traits, the Sciapodinae of the greater Washington, DC area have received little attention. Overall, the sciapodine distributions in the Nearctic region are still poorly understood. Our current knowledge of the sciapodine species composition of the greater Washington, DC area is drawn largely from Robinson's (1960, 1964) review of the dolichopodid flies of the southeastern United States in which he provided state records but limited information on local distributions. In addition, two biodiversity publications have listed taxa collected from the Potomac Gorge (Brown 2008, Evans 2008). The present study aims to provide an updated inventory of the Sciapodinae of the greater Washington, DC area that may serve as a baseline for future biodiversity comparison.

METHODS

A list of sciapodine species of the greater Washington, DC area was generated based on an examination of specimens in the collection of the National Museum of Natural History (USNM), Washington, DC, USA, which is known to have significant holdings from the study area, augmented by collections of the author. Flies were identified to species using Van Duzee (1915), Curran (1942), Robinson (1960, 1964), Steyskal (1966, 1973), and other reference material. Voucher specimens from the author are deposited at USNM.

The geographic scope of the greater Washington, DC area is defined for the purpose of this faunal paper as the surrounding area approximately 65 km (40 mi) from the center of Washington, DC. A collections locality map was produced using SimpleMappr software (Shorthouse 2010). Latitude and longitude coordinates were recorded from specimen labels. If coordinates were unavailable, latitude and longitude were estimated from maps to determine the geographical coordinates. Collection sites included the Chesapeake and Ohio Canal National Historical Park (MD), Difficult Run (VA), Great Falls Park (VA), Four Mile Run Park (VA), Holmes Run Scenic Easement (VA), Plummers Island (MD), Kenilworth Park and Aquatic Gardens (DC), and Rock Creek Park (DC), in addition to other natural and residential areas.

RESULTS and DISCUSSION

A total of 1657 specimens of sciapodine flies were identified from this region comprising 19 species in four genera (Table 1), representing 27% of the total Nearctic Sciapodinae diversity. The sciapodine species richness in the greater Washington, DC area was estimated using the nonparametric Chao-2 method in the program EstimateS, Version 9.1.0 (Chao 1987, Colwell 2016). Chao-2 provided an estimate of 19.25 sciapodine species for the region [the observed number of species (S_{obs}) is 19, the number of unique collections (Q_1) is 2, and the number of duplicates (Q_2) is 3]. The 95% confidence interval on the richness estimate ranged from 19.01 to 23.78 species. While the Chao-2 result of 19.25 versus the observed 19 species indicates a reasonably complete sampling of the region, the confidence interval of 19.01 to 23.78 species suggests there may be additional taxa.

Based on label data, there were 342 individual collection events. Approximately 73% ($n = 1208$) of the specimens were collected by Malaise trap; 5% ($n = 87$) by black light or

light trap; 2% (n = 28) by yellow pan trap; 1% (n = 9) by hand collecting; and 20% (n = 325) by unknown method. The distribution of all collection localities in the greater Washington, DC area is shown in Figure 1. Flies were recorded from 64 sites, including 52% (n = 33) from Maryland, 34% (n = 22) from Virginia, and 14% (n = 9) from Washington, DC. Figure 1 displays just the collection records in and around the study area; species recorded here are found throughout the eastern United States and not necessarily restricted to the Washington, DC area.

Table 1. Species of Sciapodinae (Diptera: Dolichopodidae) from the greater Washington, DC area.

Genera/Species	District of Columbia	Maryland	Virginia	Total Number of Flies Identified
AMBLYSILOPUS Bigot				
<i>Amblysilopus dorsalis</i> (Loew)	1	1*		2
<i>Amblysilopus scintillans</i> (Loew)	1	32	3	36
<i>Amblysilopus unicoiensis</i> (Robinson)			11	11
<i>Amblysilopus unifasciatus</i> (Say)		4		4
CONDYLOSTYLUS Bigot				
<i>Condylostylus banksii</i> (Van Duzee)		3*	5	8
<i>Condylostylus calcaratus</i> (Loew)		13	5	18
<i>Condylostylus caudatus</i> (Wiedemann)	8	117	928	1053
<i>Condylostylus comatus</i> (Loew)	9	23	17	49
<i>Condylostylus flavipes</i> (Aldrich)		5		5
<i>Condylostylus inermis</i> (Loew)	2	1		3
<i>Condylostylus nigrofemoratus</i> (Walker)	1	26	33	60
<i>Condylostylus patibulatus</i> (Say)	5	41	136	182
<i>Condylostylus scaber</i> (Loew)		12	12	24
<i>Condylostylus siphon</i> (Say)	5	47	84	136
<i>Condylostylus viridicoxa</i> (Aldrich)	3	6	28	37
MESORHAGA Schiner				
<i>Mesorhaga albiciliata</i> (Aldrich)		1*		1
SCIAPUS Zeller				
<i>Sciapus filipes</i> (Loew)		4	20	24
<i>Sciapus pallens</i> (Wiedemann)	1			1
<i>Sciapus tener</i> (Loew)		2	1	3
Total Number of Flies Identified	36	338	1283	1657
Total Number of Species	10	17	13	19

*New state record.




Figure 1. Map of the greater Washington, DC area displaying distribution of collection records of Sciapodinae (Diptera: Dolichopodidae), with eastern USA (inset). The map was generated using SimpleMappr software (Shorthouse 2010) applying the World Mercator map projection. The presented map displays just the collection records in and around the greater Washington, DC area.

There was considerable overlap of species recorded during the present study compared with previously published records (Robinson 1960, 1964; Brown 2008; Evans 2008). All taxa previously recorded from the Washington, DC area were recorded in the present inventory. Herein, the species *Amblypsilopus dorsalis* (Loew) and *Condylostylus banksii* (Van Duzee) are reported for the first time in Maryland (Table 1, Figure 2). Robinson (1960) recorded *A. dorsalis* from Rock Creek Park in Washington, DC and its known distribution ranges south into Florida and the Neotropics (Pollet et al. 2004). Presently, few localities exist for *C. banksii*; it is currently reported from Falls Church, Virginia, in addition to Massachusetts and New York (Pollet et al. 2004). Also notable was the presence of *Mesorhaga albiciliata* (Aldrich) (Table 1, Figure 2) in Maryland, which is only known from Georgia, New Jersey, and North Carolina (Robinson 1964, Pollet et al. 2004).



Figure 2. Sciapodinae species new to Maryland. (a) *Amblypsilopus dorsalis* (Loew), lateral view of female; (b) *Condyllostylus banksii* (Van Duzee), lateral view of male; (c) *Mesorhaga albiciliata* (Aldrich), lateral view of female. Photographed by Gary D. Ouellette.

The most commonly collected sciapodine species in this geographic area, by total occurrences, included *Condylostylus caudatus* (Wiedemann) (64% of all specimens, n = 1053), *C. patibulatus* (Say) (11%, n = 182), and *C. siphon* (Say) (8%, n = 136), which collectively comprised approximately 83% (n = 1371) of all identified specimens. All three species are common and widely distributed throughout much of North America (Curran 1942, Pollet et al. 2004). Overall, the genus *Condylostylus* Bigot represented 95% (n = 1575) of all identified sciapodine specimens. However, this large, diverse genus is complex and the taxonomy of a few of the recorded species are questionable (Robinson 1964). Both *C. calcaratus* (Loew) and *C. inermis* (Loew) are considered variants of *C. nigrofemoratus* (Walker) that differ only in male secondary features and probably do not represent true species (Robinson, in litt., 4 May 2019).

Robinson (in litt.) further elaborated that dolichopodids are much rarer in the area than they were in the past as a possible result of pesticide spraying and other human alterations to the environment.

Although it is possible that some sciapodine species were not represented in the samples examined, the present investigation serves as baseline to support future ecological and biodiversity studies for the greater Washington, DC area and surrounding environs.

ACKNOWLEDGMENTS

The author gratefully thanks Justin B. Runyon (United States Department of Agriculture, Forest Service) and Jung W. “Woogie” Kim (United States Department of Agriculture, Animal and Plant Health Inspection Service), as well as, Harold Robinson and John W. Brown (National Museum of Natural History) for helpful comments on the manuscript. The author also acknowledges Norman E. Woodley and F. Christian Thompson for facilitating study of material in the USNM collections.

LITERATURE CITED

- Alexander, C.P., and W.L. McAtee. 1920. Diptera of the superfamily Tipuloidea found in the District of Columbia. *Proceedings of the United States National Museum* 58(2344):385–435.
- Banks, N., C.T. Greene, W.L. McAtee, and R.C. Shannon. 1916. District of Columbia Diptera: Syrphidae. *Proceedings of the Biological Society of Washington* 29:173–204.
- Bickel, D.J. 1994. The Australian Sciapodinae (Diptera: Dolichopodidae), with a review of the Oriental and Australasian faunas, and a world conspectus of the subfamily. *Records of the Australian Museum, Supplement* 21:1–394.
- Brown, J.W. 2008. Appendix: List of the invertebrates of Plummers Island, Maryland. *Bulletin of the Biological Society of Washington* 15(1):192–226.

- Brown, J.W., and S.M. Bahr, II. 2008. The insect (Insecta) fauna of Plummers Island, Maryland: Brief collecting history and status of the inventory. *Bulletin of the Biological Society of Washington* 15(1):54–64.
- Chao, A. 1987. Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* 43(4):783–791.
- Colwell, R.K. 2016. EstimateS: Statistical estimation of species richness and shared species from samples, Version 9.1.0. User's guide and application published at: <http://viceroy.eeb.uconn.edu/estimates>.
- Curran, C.H. 1942. American Diptera. *Bulletin of the American Museum of Natural History* 80(3):51–84.
- Evans, A.V. (Coordinator). 2008. The 2006 Potomac Gorge BioBlitz: Overview and results of a 30-hour rapid biological survey. *Banisteria* 32:3–80.
- Green, M.J.B., R. How, U.K.G.K. Padmalal, and S.R.B. Dissanayake. 2009. The importance of monitoring biological diversity and its application in Sri Lanka. *Tropical Ecology* 50(1):41–56.
- Hughes, J.B., G.C. Daily, and P.R. Ehrlich. 2000. Conservation of insect diversity: A habitat approach. *Conservation Biology* 14(6):1788–1797.
- Longino, J.T., and R.K. Colwell. 1997. Biodiversity assessment using structured inventory: Capturing the ant fauna of a tropical rain forest. *Ecological Applications* 7(4):1263–1277.
- Malloch, J.R., C.T. Greene, and W.L. McAtee. 1931. District of Columbia Diptera: Rhagionidae. *Proceedings of the Entomological Society of Washington* 33(9):213–220.
- Mathis, W.N., and G.A. Foster. 2007. Canacidae (Diptera) from the Delmarva States. *Proceedings of the Biological Society of Washington* 120(4):387–428.
- McAtee, W.L. 1921. District of Columbia Diptera: Scatopsidae. *Proceedings of the Entomological Society of Washington* 23(5):120–124.
- Perry, M.C. (Editor). 2007. *The Washington Biologists' Field Club: Its members and its history (1900–2006)*. Published by The Washington Biologists' Field Club, Washington, DC. 342 pp.
- Pollet, M. 2001. Dolichopodid biodiversity and site quality assessment of reed marshes and grasslands in Belgium (Diptera: Dolichopodidae). *Journal of Insect Conservation* 5(2):99–116.
- Pollet, M., and P. Grootaert. 1991. Horizontal and vertical distribution of Dolichopodidae (Diptera) in a woodland ecosystem. *Journal of Natural History* 25(5):1297–1312.

- Pollet, M., and P. Grootaert. 1996. An estimation of the natural value of dune habitats using Empidoidea (Diptera). *Biodiversity and Conservation* 5(7):859–880.
- Pollet, M., H. Meuffels, and P. Grootaert. 2003. Status of Dolichopodidae of the Flemish Red Data Book in the Netherlands (Insecta: Diptera). Pages 61–68 in: M. Reemer, P.J. van Helsdingen, and R.M.J.C. Kleukers (Editors), *Changes in Ranges: invertebrates on the move. Proceedings 13th International Colloquium European Invertebrate Survey, Leiden, 2–5 September 2001*. European Invertebrate Survey – The Netherlands, Leiden. 137 pp.
- Pollet, M.A.A., S.E. Brooks, and J.M. Cumming. 2004. Catalog of the Dolichopodidae (Diptera) of America North of Mexico. *Bulletin of the American Museum of Natural History* 283:1–114.
- Robinson, H. 1960. Four new Dolichopodidae from the eastern United States (Diptera). *The Ohio Journal of Science* 60(5):271–273.
- Robinson, H. 1964. A synopsis of the Dolichopodidae (Diptera) of the southeastern United States and adjacent regions. *Miscellaneous Publications of the Entomological Society of America* 4(4):103–192.
- Shorthouse, D.P. 2010. SimpleMappr, an online tool to produce publication-quality point maps. Available at: <http://www.simplemappr.net>. Accessed 30 March 2019.
- Steyskal, G.C. 1966. A new species of *Sciapus* from California, with a revised key to the species of *Sciapus* found in America north of Mexico (Diptera, Dolichopodidae). *Proceedings of the Entomological Society of Washington* 68(4):290–294.
- Steyskal, G.C. 1973. Notes on southeastern North American species of *Condylostylus* and *Sciapus* (Diptera: Dolichopodidae). *The Florida Entomologist* 56(4):359–360.
- Van Duzee, M.C. 1915. Descriptions of three new species of dipterous genus *Sciapus* with a key to the North American species. *Entomological News* 26(1):17–26.

**Yellow-banded Bumble Bee, *Bombus terricola* Kirby (Hymenoptera: Apidae),
Found in West Virginia**

Mark J. Hepner and Ernest W. Smith

AllStar Ecology LLC, 1582 Meadowdale Road Fairmont, West Virginia 26554
(MJH) *Mark@AllStarEcology.com*; (EWS) *Ernie@AllStarEcology.com*

In eastern North America, several bumble bee species (Hymenoptera: Apidae: *Bombus* Latreille spp.) have declined (Colla and Parker 2008; Evans et al. 2008; Grixti et al. 2009; Cameron et al. 2011, 2016; Schweitzer et al. 2012; Williams et al. 2014). One of those in decline is the Yellow-banded Bumble Bee, *Bombus terricola* Kirby (Figure 1) (Evans et al. 2008, Cameron et al. 2011, Jacobson et al. 2018). In 2018, a survey of West Virginia (WV) native bee species was initiated to determine status and distributions throughout the state. Collectors for the survey utilized Trimble Geo 7X sub-meter accuracy GPS units with customized data input to document survey locations, habitat, and other resources available at the time of surveys. Past collection records indicated that the Yellow-banded Bumble Bee was historically present within WV. The Yellow-banded Bumble Bee was previously documented in Pocahontas County in 1958, 1960, and 1972, and Pendleton County in 1960 and 1979 (Milliron 1971, McKinney 2016, Ascher and Pickering 2018, Various Contributors 2019).

Williams et al. (2014) indicate various characters for identifying the Yellow-banded Bumble Bee and for distinguishing it from other bumble bee species. The Yellow-banded Bumble Bee has short, even hair; a short head with an oculo-malar area (cheek) barely shorter than broad (differing from *Bombus pensylvanicus* (De Geer), *B. auricomus* (Robertson), *B. nevadensis* Cresson); the midleg basitarsus with a rounded rear angle; and the flat outer surface of the hindleg tibia without long hair but having long fringes along the sides forming a corbicula. The hair on the head is black or with a minimal number of short pale hairs intermixed; the base of T2 is usually yellow without black or with only a narrow fringe along the basal margin; if T2 is more extensively black, then T4–5 are also mostly black (differing from *B. occidentalis* Greene). T3 is usually yellow and T5 is black or yellow-brown (differing from most *B. cryptarum* (Fabricius)). The wings are slightly brown (differing from *B. cryptarum*) (Williams et al. 2014). The Yellow-banded Bumble Bee has a more constant color pattern as compared to other bumble bees but has a marked chromatic variation, particularly between specimens collected from the southeastern and northwestern areas of its range (Milliron 1971).

During the 2018 WV native bee survey, nine Yellow-banded Bumble Bees were found at eight locations from 14 July to 27 August. Seven females and two males were found during surveys within Pendleton, Pocahontas, Randolph, and Tucker Counties (Table 1, Figure 2). Eight of the individuals were either not specifically seen during capture or misidentified in the field. A queen captured on 27 August was field identified, photographed, and released. Table 1 lists the floral resource from which bees were



Figure 1. Female Yellow-banded Bumble Bee (*Bombus terricola* Kirby). Top: lateral view; bottom: posterior view. Collected with a hand net during a 10-minute roadside survey in Pocahontas County, West Virginia on 11 August 2018. Photographed by Mark J. Hepner.

collected during these surveys. These floral resources included sunflower, *Helianthus* L. spp.; goldenrod, *Solidago* L. spp.; thistle, *Cirsium* Mill. spp.; and wingstem, *Verbesina alternifolia* (L.) Britton ex Kearney (all Asteraceae). The most northern and most southern survey locations were 116 km (72 mi) apart, averaging a distance of 16 km (10 mi) between individual collection locations. All *B. terricola* specimens collected were in high-elevation habitat of WV (Byers et al. 2007), averaging 1,200 m (3,937 ft) above sea level and all *B. terricola* specimens were collected within the United States Department of Agriculture (USDA) United States Forest Service’s (USFS) Monongahela National Forest. The 2018 records represent the first *B. terricola* to be found in WV since 1979 (Ascher and Pickering 2018; United States Fish and Wildlife Service, in litt.). Bee identification was validated by Sam Droege at the United States Geological Survey (USGS) Patuxent Wildlife Research Center’s (PWRC) Bee Inventory and Monitoring Lab (BIML) in Laurel, Maryland.

Table 1: 2018 West Virginia Yellow-banded Bumble Bee locations.

Date Collected	County	Number Collected	Sex	Elevation - m (ft)	Latitude	Longitude	Habitat	Floral Resource
14 JUL 2018	Pocahontas	1	F	1,382 (4,534)	38.2686	-80.2393	early-successional, roadside	<i>Erigeron strigosus</i> Muhl. ex Willd., <i>Securigera varia</i> (L.) Lassen
10 AUG 2018	Pocahontas	1	F	860 (2,822)	38.3893	-80.1293	early-successional, roadside	<i>Verbesina alternifolia</i> (L.) Britton ex Kearney
11 AUG 2018	Pocahontas	2	F	1,361 (4,465)	38.5146	-79.9116	early-successional, roadside	<i>Eutrochium purpureum</i> (L.) E.E. Lamont, <i>Prunella vulgaris</i> L.
11 AUG 2018	Pendleton	1	M	1,193 (3,914)	38.6462	-79.6270	early-successional, roadside	<i>Asclepias syriaca</i> L., <i>Silphium perfoliatum</i> L., <i>Ratibida pinnata</i> (Vent.) Barnhart
11 AUG 2018	Pendleton	1	M	1,423 (4,669)	38.6925	-79.5431	early-successional, roadside	<i>Centaurea stoebe</i> L. ssp. <i>micranthos</i> (Gugler) Hayek, <i>Cirsium</i> Mill. spp.
21 AUG 2018	Randolph	1	F	1,207 (3,960)	38.7374	-79.6018	early-successional, field	<i>Helianthus</i> L. spp.
26 AUG 2018	Tucker	1	F	1,169 (3,835)	38.9679	-79.3447	early-successional, roadside	<i>Cirsium</i> Mill. spp., <i>Monarda fistulosa</i> L., <i>Solidago</i> L. spp.
27 AUG 2018	Tucker	1	F	1,002 (3,287)	39.0160	-79.3194	early-successional, roadside	<i>Solidago</i> L. spp.




Figure 2. West Virginia Yellow-banded Bumble Bee locations, 14 July–27 August 2018. Nine records, eight locations. Created with SimpleMappr (Shorthouse 2010).

ACKNOWLEDGMENTS

Thanks to Sam Droege (USGS-PWRC-BIML) for identifying collected bees and for teaching us how to collect, process, and identify bumble bees. Without the USDA-USFS-Monongahela National Forest, West Virginia Division of Natural Resources, West Virginia State Parks, and private landowners allowing surveys on lands they own or manage the project would not have been a success. Thanks to Justin DeVault and Aaron Nemeyer with AllStar Ecology LLC for participating in the 2018 WV native bee survey and helping with the identification of floral resources recorded during the surveys. The support of AllStar Ecology LLC was instrumental in the birth of this project. Thanks to Sam Droege and Jonathan Mawdsley (Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, District of Columbia) for reviewing the manuscript and for providing helpful comments.

LITERATURE CITED

- Ascher, J.S., and J. Pickering. 2018. Discover Life bee species guide and world checklist Hymenoptera: Apoidea: Anthophila). Available at: https://www.discoverlife.org/mp/20q?guide=Apoidea_species.
- Byers, E.A., J.P. Vanderhorst, B.P. Streets. 2007. *Classification and Conservation Assessment of High Elevation Wetland Communities in the Allegheny Mountains of West Virginia*. West Virginia Natural Heritage Program, Wildlife Resources Section, West Virginia Division of Natural Resources, Elkins, WV. 92 pp. + appendices A–I. Available at: <http://wvdnr.gov/publications/PDFFiles/High%20Allegheny%20Wetlands-web.pdf>.

- Cameron, S.A., J.D. Lozier, J.P. Strange, J.B. Koch, N. Cordes, L.F. Solter, and T.L. Griswold. 2011. Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences* 108(2):662–667.
- Cameron, S.A., H.C. Lim, J.D. Lozier, M. Duennes, and R. Thorp. 2016. Test of the invasive pathogen hypothesis of bumble bee decline in North America. *Proceedings of the National Academy of Sciences* 113(16):4386–4391.
- Colla, S.R., and L. Parker. 2008. Evidence for decline in eastern North American bumblebees (Hymenoptera: Apidae), with special focus on *Bombus affinis* Cresson. *Biodiversity and Conservation* 17(6):1379–1391.
- Evans, E., R. Thorp, S. Jepsen, and S.H. Black. 2008. *Status Review of Three Formerly Common Species of Bumble Bee in the Subgenus Bombus*. 63 pp. Available at: https://www.xerces.org/wp-content/uploads/2009/03/xerces_2008_bombus_status_review.pdf.
- Grixti, J.C., L.T. Wong, S.A. Cameron, and C. Favret. 2009. Decline of bumble bees (*Bombus*) in the North American Midwest. *Biological Conservation* 142(1):75–84.
- Jacobson, M.M., E.M. Tucker, M.E. Mathiasson, and S.M. Rehan. 2018. Decline of bumble bees in northeastern North America, with special focus on *Bombus terricola*. *Biological Conservation* 217:437–445.
- McKinney M. 2016. *Bee Natural History, Diversity, and Management in West Virginia*. Dissertation submitted to the Davis College of Agriculture, Natural Resources, and Design at West Virginia University, Morgantown, West Virginia. 126 pp.
- Milliron, H.E. 1971. A monograph of the western hemisphere bumblebees [sic] (Hymenoptera: Apidae; Bombinae): I. The genera *Bombus* and *Megabombus* subgenus *Bombias*. *Memoirs of the Entomological Society of Canada* 82:1–80.
- Schweitzer, D.F., N.A. Capuano, B.E. Young, and S.R. Colla. 2012. *Conservation and Management of North American Bumble Bees*. NatureServe, Arlington, Virginia, and United States Department of Agriculture Forest Service, Washington, DC. 16 pp. Available at: https://www.fs.fed.us/wildflowers/pollinators/documents/ConsMgmtNA_BumbleBees.pdf.
- Shorthouse, D.P. 2010. SimpleMappr, an online tool to produce publication-quality point maps. Available at: <http://www.simplemappr.net>. Accessed 5 August 2019.
- Various Contributors. 2019. Hymenoptera Online (HOL), Specimen OSUC 100911, *Bombus terricola* Kirby, Pendleton County, West Virginia. Available at: <https://hol.osu.edu/spmInfo.html?id=OSUC%20100911>. Accessed 25 May 2019.
- Williams, P., R. Thorp, L. Richardson, and S. Colla. 2014. *Bumble Bees of North America: An Identification Guide*. Princeton University Press, Princeton, NJ. 208 pp.

Bumble Bees, *Bombus* Latreille (Hymenoptera: Apidae), in State Wildlife Action Plans: New Opportunities for Conservation

Jonathan R. Mawdsley^{1*}, Davia M. Palmeri², and Mark Humpert³

¹*Department of Entomology, National Museum of Natural History, Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012, USA; mawdsleyj@si.edu*

**Corresponding Author*

²*Oregon Department of Fish and Wildlife, 4034 Fairview Industrial Drive SE, Salem, OR 97302, USA; Davia.M.Palmeri@state.or.us*

³*Association of Fish & Wildlife Agencies, 1100 First Street NE, Suite 825, Washington, DC 20002, USA; mhumpert@fishwildlife.org*

Abstract: We report significant growth in interest and available resources for the conservation of bumble bees, *Bombus* Latreille (Hymenoptera: Apidae), in the USA, as measured by the inclusion of these species in the 56 U.S. State Wildlife Action Plans. In the first editions of these plans, completed in 2005, only three states included a total of three species of the genus *Bombus* in their plans. In the second editions of these plans, completed in 2015–2016, 26 states and the District of Columbia included 25 species of the genus *Bombus* in their plans as “Species of Greatest Conservation Need.” The species most frequently identified by states as “Species of Greatest Conservation Need” included *Bombus affinis* Cresson (17 states and the District of Columbia), *B. pensylvanicus* (De Geer) (17 states), *B. terricola* Kirby (15 states), and *B. fervidus* (Fabricius) (11 states). A complete list of the *Bombus* species included in these plans and a map showing the associated states are provided. Inclusion of these species in these plans will increase the available funding for bumble bee conservation and provide new opportunities for interstate and regional partnerships to conserve these species.

Keywords: Apidae, Apoidea, *Bombus*, bumble bee, conservation, Hymenoptera, pollinator

Significant declines have been reported in populations of many North American species of bumble bees, *Bombus* Latreille (Hymenoptera: Apidae; Figure 1) (Brown 2011, Cameron et al. 2011, The Xerces Society for Invertebrate Conservation 2019), and multiple species in this genus have been formally proposed for listing under the U.S. Endangered Species Act (The Xerces Society for Invertebrate Conservation 2010, 2013). Reports of bumble bee population declines in the USA have generated considerable interest among conservation biologists (Brown 2011) and wildlife managers (Learn 2016). Here we document the growth of interest and enhanced opportunities for bumble

bee conservation in the USA between 2005 and 2016, as reflected by increased inclusion of *Bombus* species in the 56 U.S. State Wildlife Action Plans.



Figure 1: *Bombus auricomus* (Robertson) visiting wild bergamot, *Monarda fistulosa* L. (Lamiaceae), on the National Mall in Washington, DC, USA. This bumble bee species has been identified as a “Species of Greatest Conservation Need” in Delaware and Maryland.

The State Wildlife Action Plans (SWAPs) are documents that describe approaches for the conservation of wildlife species and ecological communities in each of the 50 U.S. states, the District of Columbia, and five U.S. territories (Riexinger and Williamson 2009, Stoms et al. 2010, Meretsky et al. 2012). Under the U.S. federal system, the governments of these individual states and territories have legal responsibility for managing much of the nation’s biodiversity, including many of the animal pollinator species which are not listed under the federal Endangered Species Act (Association of Fish & Wildlife Agencies 2012, The Heinz Center 2013, Mawdsley et al. 2016).

Each of the 56 SWAPs is intended to present a comprehensive blueprint for the conservation of aquatic and terrestrial biodiversity within a particular state or territory. Each plan, developed in collaboration with multiple conservation partners, contains a set of common elements: a list of species of conservation interest, called “Species of Greatest Conservation Need” (SGCN); descriptions of the habitats occupied by these wildlife species; descriptions of threats to species and their habitats; identification of monitoring approaches, including both status and effectiveness measures; provisions for public engagement; and provisions for review and revision of the plans (Riexinger and Williamson 2009, Stoms et al. 2010, Fontaine 2011, Meretsky et al. 2012). The first set of SWAPs was completed in 2005 (Association of Fish & Wildlife Agencies 2012), and a

second, revised, set of plans has now been completed by states and territories and published online in 2015–2016 (Mawdsley et al. 2016).

Although pollinators were not specifically identified as a priority for inclusion in the original SWAPs, many states did include taxa from insect pollinator groups in their first plans. According to an analysis and review published by The Heinz Center (2013), 127 species of butterflies (Lepidoptera: Papilionoidea) and 103 species of skippers (Lepidoptera: Hesperioidea) were mentioned in 40 of the 56 original plans; 24 plans mentioned one or more native moth species (Lepidoptera); 11 plans included flies (Diptera); and 10 plans mentioned social or solitary bees (Hymenoptera: Apoidea), including a total of 31 bee taxa. Three states (Alaska, California, and Illinois) included three species of the genus *Bombus* as SGCN (*Bombus franklini* (Frison), *B. fraternus* (Smith) and *B. occidentalis* Greene) in their original State Wildlife Action Plans (The Heinz Center 2013, Mawdsley and Humpert 2016).

A revised set of SWAPs was prepared and released by the individual U.S. state fish and wildlife agencies in 2015–2016 (Mawdsley et al. 2016). Preliminary data collected from state wildlife agency staff before the final completion of these plans indicated that many of the revised plans were likely to include bumble bees and other pollinator taxa as SGCN (Mawdsley and Humpert 2016). Because these plans are directly associated with dedicated funding from federal and regional grants programs, the inclusion of species of the genus *Bombus* in these plans will create important new opportunities for bumble bee conservation efforts in North America.

METHODS

Staff from the Association of Fish & Wildlife Agencies conducted annual or semi-annual web-based surveys of the 56 SWAP coordinators between 2013 and 2016, in order to learn more about the progress of the individual plan revisions, as well as the plant and animal taxa that were likely going to be included in the revised plans. The online “Survey Monkey” platform (www.surveymonkey.com) was used to collect this information from the SWAP coordinators. Surveys conducted in May 2015 and May 2016 specifically asked about the possible inclusion in the revised plans of representatives from animal taxa that included known or likely pollinator species. The May 2015 survey asked states and territories whether they were planning to include native bees (Hymenoptera: Apoidea) in their revised plans, while the May 2016 survey asked specifically whether the individual states and territories were including bumble bees, *Bombus* spp., in the revised plans. In four cases of non-response to this question (i.e., either the state representative did not complete the survey, or left the question about bumble bees blank), the authors followed up directly with the SWAP coordinator in the non-responsive state in order to obtain information about the inclusion of bumble bees in the revised plan.

For those states that had indicated in the surveys that they would be including bumble bees in their revised plans, the authors then reviewed final copies of their revised plan documents in 2016 as posted on official state government websites (links to all 56 plans are available at: <https://www.fishwildlife.org/afwa-informs/state-wildlife-action-plans>). For each plan, the authors downloaded the relevant portion(s) of the document which

contained the lists of SGCN in .pdf or .docx or .xlsx formats. The authors then performed a comprehensive word search for the following strings of text characters: “bumble,” “bumblebee,” “bee,” and “*Bombus*.” All species of *Bombus* that had been included as SGCN in the revised plans were then listed in a Microsoft Excel spreadsheet, along with a comprehensive list of the individual states that had included each of these species as SGCN.

RESULTS

Twenty-seven of the revised SWAP documents included at least one species of the genus *Bombus* as an SGCN. One additional state (Colorado) did not include bumble bees as SGCN, but did mention The Xerces Society for Invertebrate Conservation and collaborators’ “Bumble Bee Watch” citizen science program (www.bumblebeewatch.org) in the revised plan’s chapter on monitoring of wildlife species. Twenty-five species of *Bombus* were included in total across all of the revised plans. Lists of these species with the associated states are presented in Table 1. Species most frequently identified by states as SGCN included *Bombus affinis* Cresson (17 states and the District of Columbia), *B. pensylvanicus* (De Geer) (17 states), *B. terricola* Kirby (15 states), and *B. fervidus* (Fabricius) (11 states).

Table 1: *Bombus* Latreille species included in the 2015–2016 revised U.S. State Wildlife Action Plans. States are designated by their postal abbreviations.

Species	States including the species as a “Species of Greatest Conservation Need”
<i>B. affinis</i> Cresson	CT, DC, DE, GA, MA, MD, ME, MI, MN, NC, NH, NJ, NY, PA, RI, VA, VT, WI
<i>B. ashtoni</i> (Cresson)	CT, DE, MD, ME, NJ, NY, PA, VT
<i>B. auricomus</i> (Robertson)	DE, MD
<i>B. bohemicus</i> (Seidl)	MN, VA
<i>B. borealis</i> Kirby	GA, NY
<i>B. citrinus</i> (Smith)	MD, ME, VT
<i>B. fernaldae</i> (Franklin)	ME, VT
<i>B. fervidus</i> (Fabricius)	ID, MA, ME, MN, NC, NH, NJ, NY, VA, VT, WI
<i>B. franklini</i> (Frison)	CA, OR
<i>B. fraternus</i> (Smith)	DE, IL, NC, NJ, OK, VA
<i>B. frigidus</i> Smith	WI
<i>B. griseocollis</i> (De Geer)	ME
<i>B. huntii</i> Greene	ID
<i>B. insularis</i> (Smith)	ID, WA
<i>B. morrisoni</i> Cresson	ID, WA
<i>B. occidentalis</i> Greene	AK, CA, ID, OR, WA
<i>B. pensylvanicus</i> (De Geer)	CT, DE, ID, LA, MA, MD, ME, MN, NC, NH, NJ, NY, OK, TX, VA, VT, WI
<i>B. perplexus</i> Cresson	VT, WI
<i>B. rufocinctus</i> Cresson	VT
<i>B. sandersoni</i> Franklin	MD, ME, NJ, WI
<i>B. sonorus</i> Say	TX
<i>B. suckleyi</i> Greene	ID, WA
<i>B. terricola</i> Kirby	CT, MA, MD, ME, MI, MN, NC, NH, NJ, NY, PA, RI, VA, VT, WI
<i>B. vagans</i> Smith	DE, MD, NC
<i>B. variabilis</i> (Cresson)	DE, MD, NC, NJ, TX, VA

DISCUSSION

The SWAPs are closely linked to important funding sources for wildlife conservation in the United States, particularly the State and Tribal Wildlife Grants Program which is administered by the U.S. Fish and Wildlife Service (Association of Fish & Wildlife Agencies 2011, 2012). This grant program provides each state with annual funding for the conservation of those wildlife species that are not the subject of active hunting or trapping programs (the so-called “non-game” species). Each state receives a direct apportionment of funding from this program each year, and funds are also available for competitive grant proposals to support projects that benefit multiple states. The financial support from this program is intended to benefit the conservation of species which are included as SGCN in the SWAPs (Association of Fish & Wildlife Agencies 2011, 2012).

By including bumble bees as SGCN in their revised SWAPs, 26 states and the District of Columbia now have expanded opportunities to conduct conservation activities to benefit these species. One of the most important sources of financial support for these conservation activities is the State and Tribal Wildlife Grants Program, which has contributed over one billion US\$ towards the conservation of SGCN and their habitats since the program’s inception in the year 2000 (Association of Fish & Wildlife Agencies 2012). Activities that could potentially be funded through these grants program include surveys and monitoring for rare bumble bee species, status reviews and the development of conservation plans for individual species or groups of species, and projects to restore and enhance bumble bee habitats. In addition, multiple states could work together to develop broader, cross-boundary conservation strategies for rare and declining bumble bee species using dedicated funding available through the competitive portion of the State and Tribal Wildlife Grants Program. Finally, funding may also be available to conserve these species from other grant programs, such as the Northeast Association of Fish and Wildlife Agencies’ Regional Conservation Needs Grant Program (Northeast Fish and Wildlife Diversity Technical Committee 2015). Together, these resources offer significantly expanded opportunities for bumble bee conservation in the United States.

ACKNOWLEDGMENTS

We thank the Wildlife Diversity Program Managers and State Wildlife Action Plan Coordinators in the 56 U.S. state and territorial fish and wildlife agencies for providing us with information about the inclusion of pollinator taxa (including bumble bees) in the revised U.S. State Wildlife Action Plans. We are especially grateful to Chris Burkett (Virginia Department of Game and Inland Fisheries), Jenny Dickson (Connecticut Bureau of Natural Resources), John Kanter (New Hampshire Department of Fish and Game), Eric Odell (Colorado Parks and Wildlife), and Mary Pfaffko (Georgia Wildlife Resources Division) for discussions that greatly improved our understanding of state conservation efforts to benefit pollinating insects. For help with the original review of the individual State Wildlife Action Plans, we thank Leslie Corcelli, Martha Surridge, and Kathryn Wallace of The Heinz Center, as well as Abigail L. Benson of the U.S. Geological Survey. We thank Sam Droege of the U.S. Geological Survey for review of the manuscript prior to publication.

LITERATURE CITED

- Association of Fish & Wildlife Agencies. 2011. *State & Tribal Wildlife Grants Program: 10 years of success*. Association of Fish & Wildlife Agencies, Washington, D.C. 70 pp.
- Association of Fish & Wildlife Agencies. 2012. *Best practices for State Wildlife Action Plans: Voluntary guidance to states for revision and implementation*. Association of Fish & Wildlife Agencies, Washington, DC. 66 pp.
- Brown, M.J.F. 2011. The trouble with bumblebees [sic]. *Nature* 469:169–170.
- Cameron, S.A., J.D. Lozier, J.P. Strange, J.B. Koch, N. Cordes, L.F. Solter, and T.L. Griswold. 2011. Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences of the United States of America* 108(2):662–667.
- Fontaine, J.J. 2011. Improving our legacy: Incorporation of adaptive management into State Wildlife Action Plans. *Journal of Environmental Management* 92(5):1403–1408.
- The Heinz Center. 2013. *Pollinators and the State Wildlife Action Plans: Voluntary guidance for state wildlife agencies*. The Heinz Center, Washington, DC. 20 pp.
- Learn, J.R. 2016. Pollinators in decline: Researchers focus on wild bees. *The Wildlife Professional* 10(2):16–22.
- Mawdsley, J.R., and M. Humpert. 2016. Revised State Wildlife Action Plans offer new opportunities for pollinator conservation in USA. *Natural Areas Journal* 36(4):453–457.
- Mawdsley, J.R., M. Humpert, and M. Pfaffko. 2016. The 2015 State Wildlife Action Plans: Meeting today's challenges in wildlife conservation. *The Wildlife Professional* 10(3):16–19.
- Meretsky, V.J., L.A. Maguire, F.W. Davis, D.M. Stoms, J.M. Scott, D. Figg, D.D. Goble, B. Griffith, S.E. Henke, J. Vaughn, and S.L. Yaffee. 2012. A state-based national network for effective wildlife conservation. *BioScience* 62(11):970–976.
- Northeast Fish and Wildlife Diversity Technical Committee. 2015. *Taking Action Together: Northeast regional synthesis for State Wildlife Action Plans*. Northeast Association of Fish & Wildlife Agencies, Dover, Delaware. 222 pp.
- Riexinger, P., and S.J. Williamson. 2009. Using State Wildlife Action Plans to guide landscape-level conservation in the northeastern United States. Pages 82–85 in R.E. McCabe and K.A. Stockwell (Editors), *Transactions of the Seventy-fourth North American Wildlife and Natural Resources Conference*. Wildlife Management Institute, Washington, DC.

Stoms, D., F. Davis, and J.M. Scott. 2010. Implementation of State Wildlife Action Plans: conservation impacts, challenges and enabling mechanisms. *Gap Analysis Bulletin* 17:30–32.

United States Geological Survey. 2019. State Wildlife Action Plans (SWAP): A national look at Species of Greatest Conservation Need as reported in State Wildlife Action Plans. Available at: <https://www1.usgs.gov/csas/swap/index.html>. Accessed 1 August 2019.

The Xerces Society for Invertebrate Conservation. 2013. *Petition to list the Rusty Patched Bumble Bee, Bombus affinis (Cresson), 1863, as an endangered species under the U.S. Endangered Species Act*. The Xerces Society for Invertebrate Conservation, Portland, Oregon. 42 pp.

The Xerces Society for Invertebrate Conservation. 2019. Red List of Bees: Native Bees in Decline. Available at: <https://xerces.org/pollinator-redlist/>. Accessed 30 July 2019.

The Xerces Society for Invertebrate Conservation and R. Thorpe. 2010. *Petition to list Franklin's Bumble Bee, Bombus franklini (Frison), 1921, as an endangered species under the U.S. Endangered Species Act*. The Xerces Society for Invertebrate Conservation, Portland, Oregon. 40 pp.

THE MARYLAND ENTOMOLOGIST

NOTICE TO CONTRIBUTORS

Contributors should prepare manuscripts according to the following instructions.

Title: The title should be brief, concise, and pertinent.

Abstract: An abstract is required for all long articles; suggested for all biologic studies more than two (2) pages in length; but is not needed for notes, distribution reports, or short observations (especially if two pages or shorter in length). The abstract should provide a capsule description of the main thrust, methods, and essential findings of the article. It should contain the scientific name, including species author(s), of the main subject species for most biological studies.

Text: Manuscripts should be submitted in MS Word™ by e-mail or on a CD. Please identify respective file name(s) for text, figure titles, and descriptions of graphs or figures. First mention of a biological organism, in the abstract and text should include the full scientific name, including species author(s). Carefully check the spelling of all scientific names. All genera and species should be in italics. Capitalize “official” common names for faunal species. Short articles and general notes (20 pages or less) are preferred. Longer manuscripts may be assessed page charges. Color “copy ready” illustrations, pictures, or digital images are preferred.

References: References should be given in an author-date format: (Lynch 1987); (Lynch 1987, 1988); (Lynch, in press); (Lynch, in litt.); (Lynch, pers. comm.); (Lynch and Lynch 1989); and (Lynch et al. 1990) for 3 or more authors. Provide evidence of acceptance for works “in press,” or cite as “unpublished”, “in litt.” (written), or “pers. comm.” (verbal); written permission is suggested as well. Citations shall be listed alphabetically, under LITERATURE CITED, as follows: **Articles:** Lynch, J.F. 1987. An annotated checklist and key to the species of ants (Hymenoptera: Formicidae) of the Chesapeake Bay region. *The Maryland Naturalist* 31(3–4):61–105. Do not abbreviate the titles of journals. **Books:** Fisher, B.L., and S.P. Cover. 2007. *Ants of North America: A guide to the genera*. University of California Press, Berkeley and Los Angeles, CA. 194 pp. **Internet:** Entomological Society of America. 2011. Common names of insects database. Available at: <http://entsoc.org/common-names>. Accessed 11 July 2011.

Tables: Tables, graphs, and line drawings should be created electronically in black and white. Color should only be used when absolutely necessary for clarity.

Illustrations: Photographs or high-definition images may be accepted if necessary or desired by the author(s) to support the text. Reproduction of photos or images may increase printing costs and authors will be expected to pay any extra charges. Photographs should be submitted in color. Figure numbers, as cited in the text, and figure legends should be keyed to each respective photograph.

The Maryland Entomologist is published annually by the Maryland Entomological Society. There are four numbers per volume. Original articles or reports on geographic or temporal distribution (primarily pertaining to Maryland and the Mid-Atlantic region), ecology, biology, morphology, genetics, systematics, behavior, etc., are welcome. Notes on distribution, behavior, occurrence, migration, life history, and other biological topics will be published. All submissions are subject to editorial review and acceptance. Articles will be peer-reviewed. E-mail submissions to Editor Eugene J. Scarpulla at ejscarp@comcast.net.

CONTENTS

Editor’s Note.....1

Incidental Capture of *Calosoma externum* (Say) (Coleoptera: Carabidae) at Various Sites across the Coastal Plain of Maryland between 1988 and 1993 during a Monitoring and Control Program Targeting Lepidopteran Agricultural Pests
Frank G. Guarnieri.....2

Beetles (Coleoptera) Associated with Flowers of Lizard’s Tail, *Saururus cernuus* L. (Saururaceae), in Calvert County, Maryland
Brent W. Steury9

Diversity of Sciapodinae (Diptera: Dolichopodidae) in the Greater Washington, District of Columbia Area
Gary D. Ouellette.....15

Yellow-banded Bumble Bee, *Bombus terricola* Kirby (Hymenoptera: Apidae), found in West Virginia
Mark J. Hepner and Ernest W. Smith23

Bumble Bees, *Bombus* Latreille (Hymenoptera: Apidae), in State Wildlife Action Plans: New Opportunities for Conservation
Jonathan R. Mawdsley, Davia M. Palmeri, and Mark Humpert28

COVER PHOTOGRAPH

Yellow-banded Bumble Bee, *Bombus terricola* Kirby (Hymenoptera: Apidae) on common milkweed, *Asclepias syriaca* L. (Asclepiadaceae). Photographed in the Monongahela National Forest, Pendleton County, West Virginia, 25 July 2019.

Photographed by Mark J. Hepner